



THOMSON REUTERS™

PENTETATE CALCIUM TRISODIUM MICROMEDEX® DETAILED DRUG INFORMATION FOR THE CONSUMER™

Detailed Drug Information for the Consumer are in depth patient handouts that provide information on medications for FDA approved uses and some non-FDA approved uses. Documents include: an overview about the drug, phonetic pronunciation, common brand names, when the medicine should not be used, how to use it, what to do if a dose is missed, how to properly store or dispose of the medicine, drugs or foods to avoid when taking the medicine, warnings, possible side effects, and when to contact a doctor.

For more information about Thomson Reuters Micromedex, visit www.micromedex.com.

Information valid as of March 17, 2011.

The information contained in the Thomson Reuters (Healthcare) Inc. products is intended as an educational aid only. It is not intended as medical advice for individual conditions or treatment. It is not a substitute for a medical exam, nor does it replace the need for services provided by medical professionals. Talk to your doctor, nurse or pharmacist before taking any prescription or over the counter drugs (including any herbal medicines or supplements) or following any treatment or regimen. Only your doctor, nurse or pharmacist can provide you with advice on what is safe and effective for you.

The use of the Thomson Reuters (Healthcare) Inc. products is at your sole risk. These products are provided "AS IS" and "as available" for use, without warranties of any kind, either express or implied. Thomson Reuters (Healthcare) Inc. makes no representation or warranty as to the accuracy, reliability, timeliness, usefulness or completeness of any of the information contained in the products. Additionally, THOMSON REUTERS (HEALTHCARE) INC. MAKES NO REPRESENTATION OR WARRANTIES AS TO THE OPINIONS OR OTHER SERVICE OR DATA YOU MAY ACCESS, DOWNLOAD OR USE AS A RESULT OF USE OF THE THOMSON REUTERS (HEALTHCARE) INC. PRODUCTS. ALL IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE OR USE ARE HEREBY EXCLUDED. THOMSON REUTERS (HEALTHCARE) INC. DOES NOT ASSUME ANY RESPONSIBILITY OR RISK FOR YOUR USE OF THE THOMSON REUTERS (HEALTHCARE) INC. PRODUCTS.

The following information is an educational aid only. It is not intended as medical advice for individual conditions or treatments. Talk to your doctor, nurse or pharmacist before following any medical regimen to see if it is safe and effective for you.

Copyright © 2011 Thomson Reuters (Healthcare) Inc. All rights reserved. Information is for individual use only and may not be sold, redistributed or otherwise used for commercial purposes.

PENTETATE CALCIUM TRISODIUM (Intravenous route, Inhalation route) - PEN-te-tate KAL-see-um trye-SOE-dee-um

Detailed Drug Information for the Consumer™

Brand Names/Available Dosage Forms

- Some commonly used brand names are:

In the U.S.A—

- Not commercially available

†

- Not commercially available in the U.S. and Canada.

Classes

Therapeutic Class:

- Heavy Metal Chelator

Uses for This Medicine

- Pentetate Calcium Trisodium is used to remove harmful substances, such as plutonium, americium, or curium from your body.
- This medicine is available only with your doctor's prescription.

Before Using This Medicine

- In deciding to use a medicine, the risks of taking the medicine must be weighed against the good it will do. This is a decision you and your doctor will make. For this medicine, the following should be considered:

Allergies—

- Tell your doctor if you have ever had any unusual or allergic reaction to this medicine or any other medicines. Also tell your health care professional if you have any other types of allergies, such as to foods, dyes, preservatives, or animals. For non-prescription products, read the label or package ingredients carefully.

Pediatric—

- Although there is no specific information comparing use of pentetate calcium trisodium in children with use in other age groups, this medicine when given by injection is not expected to cause different side effects or problems in children than it does in adults.

Geriatric—

- Many medicines have not been studied specifically in older people. Therefore, it may not be known whether they work exactly the same way they do in younger adults or if they cause different side effects or problems in older people. There is no specific information comparing use of pentetate calcium trisodium in the elderly with use in other age groups.

Pregnancy—

The following information is an educational aid only. It is not intended as medical advice for individual conditions or treatments. Talk to your doctor, nurse or pharmacist before following any medical regimen to see if it is safe and effective for you.

Copyright © 2011 Thomson Reuters (Healthcare) Inc. All rights reserved. Information is for individual use only and may not be sold, redistributed or otherwise used for commercial purposes.

	Pregnancy Category	Explanation
All Trimesters	C	Animal studies have shown an adverse effect and there are no adequate studies in pregnant women OR no animal studies have been conducted and there are no adequate studies in pregnant women.

Lactation—

- There are no adequate studies in women for determining infant risk when using this medication during breastfeeding. Weigh the potential benefits against the potential risks before taking this medication while breastfeeding.

Interactions with Medicines—

- Although certain medicines should not be used together at all, in other cases two different medicines may be used together even if an interaction might occur. In these cases, your doctor may want to change the dose, or other precautions may be necessary. Tell your healthcare professional if you are taking any other prescription or nonprescription (over-the-counter [OTC]) medicine.

Interactions with Food/Tobacco/Alcohol—

- Certain medicines should not be used at or around the time of eating food or eating certain types of food since interactions may occur. Using alcohol or tobacco with certain medicines may also cause interactions to occur. Discuss with your healthcare professional the use of your medicine with food, alcohol, or tobacco.

Other Medical Problems—

- The presence of other medical problems may affect the use of this medicine. Make sure you tell your doctor if you have any other medical problems, especially:
 - Asthma—May be worsened by pentetate calcium trisodium given by inhalation.
 - Hemochromatosis (iron disorder)—May increase the chance for serious side effects.
 - Kidney disease—May lower the rate at which the harmful substances can be removed. Dialysis may be needed to increase removal of these harmful substances.

Proper Use of This Medicine

- Harmful substances may accumulate in your bladder. Therefore, to increase the flow of urine and decrease the time your bladder contains these harmful substances, your doctor may instruct you to drink plenty of liquids and urinate often while being treated with pentetate calcium trisodium to help eliminate the harmful substances.

Dosing—

- The dose of this medicine will be different for different patients. Follow your doctor's orders or the directions on the label. The following information includes only the average doses of this medicine. If your dose is different, do not change it unless your doctor tells you to do so.
- The amount of medicine that you take depends on the strength of the medicine. Also, the number of doses you take each day, the time allowed between doses, and the length of time you take the medicine depend on the medical problem for which you are using the medicine.
- For inhalation dosage form
 - To help with the removal of harmful substances in individuals breathing in the harmful substances
 - Adults—1 gram once a day.

The following information is an educational aid only. It is not intended as medical advice for individual conditions or treatments.

Talk to your doctor, nurse or pharmacist before following any medical regimen to see if it is safe and effective for you.

Copyright © 2011 Thomson Reuters (Healthcare) Inc. All rights reserved. Information is for individual use only and may not be sold, redistributed or otherwise used for commercial purposes.

- Children—Use and dose must be determined by your doctor.
- For parenteral dosage form
 - To help with the removal of harmful substances in individuals exposed by some other way than breathing in the substances
 - Adults—1 gram once a day.
 - Children—14 milligrams per kilogram (mg/kg) once a day.

Missed Dose—

- Call your doctor or pharmacist for instructions.

Precautions While Using This Medicine

- To prevent radiation contamination of other persons or environment:
 - Using a normal toilet instead of a urinal
 - Flushing toilet several times after each use
 - Completely cleaning up any spilled urine with a tissue and flushing it away
 - Washing hands thoroughly with soap after urinating or a bowel movement
 - Immediately laundering clothes and linens soiled with urine, feces or blood; washing them separately from other clothes
- Avoid swallowing any solid substances that may be coughed up. Dispose of solid substances in sink or toilet. Flush several times if put in toilet or flush sink by washing hands thoroughly with soap after disposal.
- Extra precaution in handling urine, feces, and substances coughed up by children to avoid additional exposure to the caregiver or to the child.
- If you are breast-feeding your baby, talk to your doctor about how long you must formula feed your baby and how to dispose of the breast milk containing harmful substances.

Side Effects of This Medicine

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention.
- Also, because of the way these medicines act on the body, there is a chance that they might cause other unwanted effects that may not occur until months or years after the medicine is used. These may include certain types of cancer, such as leukemia or bladder cancer. Discuss these possible effects with your doctor.
- *Check with your doctor immediately* if any of the following side effects occur:
 - *Incidence unknown*
 - Chest pain; cough; difficulty swallowing; dizziness; fast heartbeat; hives; itching; puffiness or swelling of the eyelids or around the eyes, face, lips or tongue; shortness of breath; skin rash; tightness in chest; unusual tiredness or weakness; wheezing
- Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side

The following information is an educational aid only. It is not intended as medical advice for individual conditions or treatments. Talk to your doctor, nurse or pharmacist before following any medical regimen to see if it is safe and effective for you.

Copyright © 2011 Thomson Reuters (Healthcare) Inc. All rights reserved. Information is for individual use only and may not be sold, redistributed or otherwise used for commercial purposes.

effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them:

- *Incidence unknown*
 - Bleeding, blistering, burning, coldness, discoloration of skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at site; blistering, crusting, irritation, itching, or reddening of skin; cracked, dry, scaly skin; diarrhea; headache; lightheadedness; metallic taste; nausea; swelling
- Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional.
- Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.